

ILLUSTRATED REVIEW

Tranexamic acid evidence and controversies: An illustrated review

Nicole Relke MD¹   | Nicholas L. J. Chornenki MD¹  |
Michelle Sholzberg MDCM, MSc, FRCPC^{2,3,4} 

¹Department of Medicine, Queen's University, Kingston, ON, Canada

²Department of Medicine, St. Michael's Hospital, University of Toronto, Toronto, ON, Canada

³Department of Laboratory Medicine & Pathobiology, St. Michael's Hospital, University of Toronto, Toronto, ON, Canada

⁴Division of Hematology, Department of Medicine, St. Michael's Hospital, University of Toronto, Toronto, ON, Canada

Correspondence

Nicole Relke, Department of Medicine, Queen's University, Kingston, ON, Canada.

Email: nrelke@qmed.ca

Handling Editor: Alisa Wolberg.

Abstract

Tranexamic acid (TXA) is an antifibrinolytic agent commonly used for the treatment or prevention of bleeding. Indications for TXA are diverse, including heavy menstrual bleeding, trauma, postpartum hemorrhage, traumatic brain injury, and surgical site bleeding. Despite decades of use and a robust body of evidence, hesitancy using TXA persists in many clinical settings. This illustrated review describes the history, pharmacology, and practical considerations of TXA use. We also describe the major landmark randomized controlled trials of TXA and their implications. Finally, we review the evidence around common controversies surrounding TXA such as the risk of thrombosis, prescription along with combined hormonal contraceptives, and use in patients with gross hematuria.

KEYWORDS

antifibrinolytic agents, blood coagulation, contraceptive agents, thrombosis, tranexamic acid

Essentials

- Tranexamic acid (TXA) decreases the risk of bleeding and often the risk of death from bleeding.
- In general, TXA does not increase the risk of blood clots.
- Consider shared decision making in patients taking TXA and combined hormonal contraceptives.
- It is unknown if TXA causes harm in patients with blood in the urine.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Research and Practice in Thrombosis and Haemostasis* published by Wiley Periodicals LLC on behalf of International Society on Thrombosis and Haemostasis (ISTH).

History of Tranexamic acid

The Okamoto Legacy¹



Left: Shosuke Okamoto, Right: Utako Okamoto

1940s

1945: Drs. Utako and Shosuke Okamoto, wife and husband team, were medical doctors and researchers at Kobe and Keio Medical School in Japan. After World War II, they directed their research towards hemostasis due to scarce resources

"If there was not enough [resources], we could simply use our own [blood]"

1960s

1960s: Post-partum hemorrhage (PPH) was identified as the major cause of maternal death in Japan. Utako and Shosuke Okamoto began to develop new compounds that could reduce the risk of PPH

1960s: Studied anti-fibrinolytic **epsilon-amino-caproic acid (EACA)**
 → Determined that a more potent agent was required

After 1962: Utako and Shosuke Okamoto were unable to persuade obstetricians to conduct research studies on the use of TXA for PPH

1962: Discovered **1-(aminomethyl)-cyclohexane-4-carboxylic acid (AMCHA)²**, a chemical relative of EACA that is 27x more powerful. AMCHA was later renamed **tranexamic acid (TXA)**

November 1, 2004: Shosuke Okamoto died

2004

2010: CRASH-2 trial showed TXA safely reduced the risk of death in bleeding trauma patients³

2010

2011: TXA added to WHO list of essential medicines⁴

2011

2014: Principle investigator of **WOMAN Trial** investigating TXA for PPH visits Utako in Japan. Utako said: "It is going to be effective"

2014

April 14, 2016: WOMAN Trial reached recruitment target of 20,000 patients

2016

April 21, 2016: Utako Okamoto died in Kobe, Japan at 98 years of age

2017

2017: WOMAN Trial showed that TXA safely decreased the risk of hemorrhagic death in women with PPH⁵


Did you know?💡

Earlier in her career, Utako worked long hours in the lab while also caring for her daughter. She was once asked to leave a conference because "events were not for women and children". Unfortunately, sexism in academia and in medicine remains prevalent today⁶


#SHERO



Results of the **WOMAN Trial** shown below!



Difference between fibrin degradation products (FDPs) and D-dimer?




FDPs can be present in the absence of stable clot formation when plasmin cleaves circulating fibrinogen

D-dimer is a FDP consisting of two D domains formed by degradation of crosslinked fibrin polymers. Indicates formation of a stable fibrin-rich clot crosslinked by **factor XIIIa**^{10,11} and subsequent fibrinolysis

Caution ⚠️

D-dimer can be elevated in thrombosis, infection (e.g. COVID-19), DIC, inflammation, malignancy, trauma, pregnancy, liver disease, older age, and recent surgery¹¹

Tranexamic Acid: Mechanism of Action



TXA Indications

The Grading of Recommendations, Assessment, Development and Evaluations (GRADE) system was used to classify quality of evidence as *high*, *moderate*, or *low*¹³

Green = High quality evidence
Yellow = Moderate quality evidence
Orange = Low quality evidence

Indication	TXA Regimen
Postpartum hemorrhage (PPH) ^{5,14} WOMAN TRIAL	1g IV over 10 min. If bleeding continues after 30 min or restarts within 24h a 2 nd dose of 1g IV can be given
Trauma-associated hemorrhage ³ CRASH-2 TRIAL	1g IV over 10 min then 1g over the next 8 hours as a continuous infusion
Reducing transfusion in cardiac surgery ¹⁵	50mg/kg IV over 30 min during the OR
Traumatic brain injury with GCS > 9 ¹⁶ CRASH-3 TRIAL	1g IV over 10 min then 1g over the next 8 hours as a continuous infusion
Heavy Menstrual Bleeding ^{17,18,19}	1300mg PO 3 times daily (3900mg/day) for up to 5 days during each monthly menstruation
Intracerebral Hemorrhage (ICH) ²⁰ TICH-2 TRIAL	1g IV over 10 min then 1g over the next 8 hours as a continuous infusion
Hemoptysis ^{21,22}	500-1000mg nebulized in 5-10mL 0.9% normal saline
Reducing transfusion during orthopedic surgery ^{23,24}	10mg/kg IV loading dose during the OR followed by 1mg/kg/hour maintenance infusion
Von Willebrand Disease (VWD) related bleeding ^{25,26}	Oral: TXA 20mg/kg PO TID Mouthwash: TXA 5% 10mL QID prn – Swish and spit
Topical surgical field blood loss reduction ²⁷	Variable doses; most commonly 1g TXA in 50 mL administered intraarticularly
Epistaxis ^{28,29}	Topical: Cotton gauze soaked with injectable form of TXA (500mg in 5mL)
Hereditary Hemorrhagic Telangiectasia related bleeding ³⁰	1g PO TID
Prophylaxis in Acute Myeloid Leukemia related bleeding ³¹	1g IV q6h when platelet count <20 or falling trend <50, until platelets >20 on two counts
Prophylaxis in Acute Promyelocytic Leukemia related bleeding ³¹	2g IV q8h for 6 days
Melasma ³³	250mg PO BID or 4mg/mL injected intradermal to melasma lesions
Hereditary angioedema ³⁴	Long-term prophylaxis: 0.5-1g PO BID-TID Short-term prophylaxis: 1g PO QID x48 before and after procedure

Did you know?



The **TRAAP study** showed that in women with **vaginal delivery** who received prophylactic oxytocin, **TXA did NOT result in significantly lower rates of measured PPH** compared to placebo³⁵

Research In Progress



Does prophylactic TXA in women with **cesarean delivery** reduce the risk of PPH?

TRAAP2 study³⁶: NCT03431805


MFMU Study: NCT03364491

WOMAN-2 → Does TXA prevent PPH in women with **vaginal delivery** and **moderate or severe anemia**?³⁷ NCT03475342

Research In Progress



Side Effects of Systemic TXA^{38,39,40}



TXA Associated Seizure

TXA is a competitive antagonist of GABA_A and glycine receptors⁴¹

Glycine TXA

TXA is a structural analogue of glycine

Did you know?

GABA_A and glycine receptors are inhibitory receptors in the CNS

↓ Inhibitory Current → ↑ Excitability → Seizure

● = TXA ● = Glycine / GABA

Risk factors^{15,41-45}

- Open-chamber cardiac surgery
- Cardio-pulmonary bypass time >150min
- High dose TXA (>80-100mg/kg total)
- Renal dysfunction
- Age >75 years
- Poor overall health

Dose-effect response of TXA-associated seizure⁴⁵

Cumulative incidence rate of TXA associated seizure⁴⁴
~2.7%*
(*cardiac surgery, pulmonary endarterectomy)

Treatment: Anesthetics, Lorazepam (↑GABA)


Caution ⚠

Urinary excretion is the main route of elimination for TXA. Dose adjustment for renal impairment required!

Serum creatinine	Adjusted TXA Dose ^{38,39}
120-250 µmol/L	15mg/kg PO or 10mg/kg IV BID
250-500 µmol/L	15mg/kg PO or 10mg/kg IV daily
>500 µmol/L	15mg/kg PO or 10mg/kg q48h

TXA Contraindications

Canada³⁸




Subarachnoid Hemorrhage

Antifibrinolytics in **aneurysmal SAH**⁴⁶:

- ✓ **Reduced risk of rebleeding** by ~35%
- ✗ **Increased risk of cerebral ischemia**



↓

No net clinical benefit




TXA Hypersensitivity⁴⁷

Mild/ Moderate: pruritis, urticaria
Severe: Anaphylaxis

Mechanism:
IgE mediated  or Cellular mediated 


Protocol exists for verification of TXA anaphylaxis⁴⁸



Hematuria



Controversy!
The use of TXA for hematuria is discussed below




Disturbances of color vision


Visual impairment, blurred vision, impaired color vision can occur³⁸

➔ **Discontinue TXA if these symptoms develop**

Consider regular ophthalmic check-up with **long-term uninterrupted TXA**




Thrombosis
Active or history of thrombosis




Controversy!
Is TXA associated with a risk of thrombosis? Discussed below


United States³⁹



Subarachnoid Hemorrhage



TXA Hypersensitivity



Thrombosis
Active or history of thrombosis
Avoid use with medications that may be pro-thrombotic: Factor IX complex concentrates, **hormonal contraceptives**

Did you know? 💡

Despite the risk of cerebral infarction, 2012 AHA guidelines for the management of **aneurysmal SAH** recommend short term (<72h) TXA or aminocaproic acid to reduce the incidence of aneurysmal rebleeding when there is an unavoidable surgical delay (**Class IIa, Level of Evidence B**)⁴⁹


⚠️ Off-label use of TXA ⚠️

What does newer evidence say (2020)

ULTRA Trial⁵⁰: In patients with **aneurysmal SAH** ➔ Ultra-early (<24h), short-term TXA did not improve clinical outcomes

✗ **Based on new evidence, routine use of TXA for aneurysmal SAH is not recommended**

Controversy!
This is discussed below.



Select Landmark TXA Randomized Controlled Trials

★ *primary outcome*

HMB¹⁹

ELIGIBILITY Adult women with **heavy menstrual bleeding** (mean menstrual blood loss (MBL) ≥ 80 mL)
Exclusion: thromboembolic disease, ocular disease, **severe anemia**, pregnant, lactating, endometrial or cervical abnormalities

METHODS **n=187**
Placebo n=72 **TXA** n=115 **TXA 1.3g PO TID x 5 days at the onset of HMB for six menstrual cycles**

RESULTS

	TXA	Placebo	P
★ Average reduction MBL (mL)	69.6	12.6	<0.001
★ Reduction MBL >50mL	56%	19%	<0.001
★ % cycles with clinically meaningful reduction in MBL (>36mL)	69%	29%	<0.001

Improvements in **health-related quality of life (HR-QoL)** in TXA group compared to placebo ($P < 0.01$)
 Adverse events were *mild/moderate in severity* (e.g. headache)

CONCLUSION TXA was **well tolerated**, and **significantly improved MBL** and **HR-QoL** in women with HMB

What is Heavy Menstrual Bleeding (HMB)?^{51,52}



Menstrual Blood Loss (MBL) >80mL

MBL that interferes with:



Physical Emotional Social
Quality of Life

More info at:
www.letstalkperiod.ca

WOMAN⁵

ELIGIBILITY Woman >16 yo with **post partum hemorrhage**

METHODS **n=20060**
TXA n=10051 **Placebo** n=10009 **TXA 1g IV Repeat if bleeding continues after 30min or rebleed within 24h**

RESULTS

	TXA	Placebo	P
★ Composite of all cause mortality or hysterectomy	5.3%	5.5%	0.65
Death from bleeding	1.5%	1.9%	0.045

No difference between groups in adverse effects

CONCLUSION TXA had no effect on all cause mortality but safely **reduced risk of death due to bleeding** in women with post partum hemorrhage

Management of HMB in women with VWD:⁵³

Who do NOT wish to conceive

★ **Hormonal therapy** (CHC or levonorgestrel-releasing intrauterine system) or **TXA** recommended over desmopressin (DDAVP)

- TXA likely reduces MBL more than DDAVP⁵⁴

Who do wish to conceive

★ **TXA** recommended over DDAVP

Research In Progress



WOMAN-2 → Does TXA prevent PPH in women with **vaginal delivery** and **moderate or severe anemia**?³⁷ NCT03475342

Select Landmark TXA Randomized Controlled Trials

★ *primary outcome*

2010

CRASH-2³

ELIGIBILITY Adult **trauma patients** within 8 hours of injury

METHODS

TXA **n=10067** **n=20211** Placebo **n=10060**

TXA 1g IV over 10 min then:
TXA 1g IV over 8 hrs

RESULTS

	TXA	Placebo	P
★ Mortality	14.5%	16.0%	0.003
Death from bleeding	4.9%	5.7%	0.007

No difference between groups in adverse events

CONCLUSION Early TXA (<3h) safely **reduced risk of death from bleeding** in trauma patients

2010

Cardiac Surgery¹⁵

ELIGIBILITY Adult patients undergoing **cardiac surgery**

METHODS

TXA **n=2329** **n=4662** Placebo **n=2333**

TXA 100mg/kg IV, protocol changed to 50mg/kg halfway through trial

RESULTS

	TXA	Placebo	P
★ Death or Thrombosis	16.7%	18.1%	0.22
Transfusion	37.9%	54.7%	<0.001
Hemorrhage + reoperation	1.4%	2.8%	<0.001
✗ Seizure	0.7%	0.1%	0.002

CONCLUSION TXA **reduced risk of transfusion & reoperation for hemorrhage**. No increased risk of death or thrombosis. TXA increased risk of seizure

2018

TICH-2²⁰

ELIGIBILITY Adults with primary **intracranial hemorrhage** within the last 8 hours

METHODS

TXA **n=1161** **n=2325** Placebo **n=1164**

TXA 1g IV over 10 min then:
TXA 1g IV over 8 hrs

RESULTS ★ **Functional status at 90 days** did not differ significantly between groups

	TXA	Placebo	P
Mortality at 7 days	9%	11%	0.04
Mortality at 90days	22%	21%	0.37

No difference between groups in adverse events

CONCLUSION TXA **did not affect functional status at 90 days**. Suggestion of decreased risk of early death

2019

CRASH-3¹⁶

ELIGIBILITY **Acute traumatic brain injury (TBI)** within 3 hours of injury

METHODS

TXA **n=4613** **n=9202** Placebo **n=4514**

TXA 1g IV over 10 min then:
TXA 1g IV over 8 hrs

RESULTS

	TXA	Placebo	RR
★ All head-injury related death (28d)	18.5%	19.8%	0.94
Severe (GCS 3-9)	39.6%	40.1%	0.99
GCS 9-15	5.8%	7.5%	0.78

No difference between groups in adverse events

CONCLUSION **Early TXA (<3h)** reduced risk of head injury related death. TXA appeared safe in TBI.
(see next page)

Select Landmark TXA Randomized Controlled Trials


★ primary outcome

2020 HALT-IT⁵⁵

ELIGIBILITY Adult with significant **upper or lower gastrointestinal (GI) bleed**


Including: patients with cirrhosis

METHODS



TXA 1g IV over 10 min then:
TXA 3g IV over 24h

RESULTS



★ Death due to bleeding (5d)
■ TXA ■ Placebo

Risk of VTE (DVT, PE) higher in patients with variceal bleeding or liver disease ($p=0.035$)

CONCLUSION

TXA **did not reduce death** from GI bleeding

- Risk of **VTE** and **seizure** higher in the TXA group

Factors Potentially Increasing VTE in HALT-IT Trial⁵⁶



↑ **Average Age** (58 yrs vs. 35 yrs in CRASH-2 trial)

↑ **Comorbidities** (7% malignancy)



Liver disease (40%):
↓ fibrinolysis in cirrhosis⁵⁷ may be associated with ↑VTE⁵⁸



Higher dose
(may also explain increased risk of seizure)


TXA not recommended for empiric treatment of GI bleeding at this time

2020

a-TREAT⁵⁹

ELIGIBILITY Patients w/ **hematologic malignancy** undergoing chemotherapy or hematopoietic stem cell transplantation with **platelets <30**

METHODS



TXA 1g IV or TXA 1.3g PO q8h for platelets <30 until platelets >30

RESULTS

	TXA	Placebo	P
★ Grade 2+ bleeding	45.4%	48.8	0.74
Central line occlusion	19.5%	11.0%	N/A*

No increase in non-catheter thrombotic events

CONCLUSION

TXA **had no effect** on the incidence of WHO Grade 2+ bleeding in this population. **Increased line occlusion** in the TXA arm but no other thrombotic events

*Awaiting full text publication


WHO bleeding scale⁶⁰

Grade 0	No bleeding
Grade 1	Petechial bleed
Grade 2	Mild blood loss (clinically significant)
Grade 3	Gross blood loss requires transfusion (severe)
Grade 4	Debilitating blood loss, retinal/cerebral bleed (associated with fatality)

Did you know?

CRASH-2, **CRASH-3** and **WOMAN** trials showed that early tranexamic acid (<3 hours) is associated with optimal effect

Mild and moderate GCS score



CRASH-3¹⁶

Early TXA (<3h)
+
 Mild-moderate head injury





=
Greatest mortality benefit

Controversy #1: TXA and Thrombosis

Given the anti-fibrinolytic effects of TXA, concern exists regarding increased risk of **thromboembolic events**

Non-surgical Patients

Risk of arterial and venous thrombosis in **non-surgical patients** receiving systemic TXA⁶¹:
 • Systematic review of 22 RCTs (including **CRASH-2** and **WOMAN**)

		 DVT	 PE	 MI	 Stroke
Number of trials (n)		8 (46630)	6 (43161)	3 (42470)	5 (42815)
Weighted event rates	TXA	0.28%	0.52%	0.27%	0.45%
	No TXA	0.29%	0.54%	0.30%	0.41%
RR (95% CI)		0.97 (0.69-1.37)	0.97 (0.75-1.26)	0.88 (0.43-1.84)	1.10 (0.68-1.78)


No increased risk of venous or arterial thrombosis in non-surgical patients receiving systemic TXA

Patients with a known history of thrombosis were excluded from these studies. In the **WOMAN** trial, patients with a known thrombotic event during pregnancy were excluded

Surgical Patients



Studies of **surgical patients** have also not shown a significant increase in thrombotic events for any surgery type while reducing blood loss^{62,63,64}

Did you know? 💡
 There was no dose response relationship for reducing surgical blood loss with TXA doses above 1g IV⁶⁴



In the absence of patient specific factors (e.g. history of thrombosis or cirrhosis) **evidence suggests there is no reason to avoid TXA in medical or surgical patients for fear of thrombosis**


Controversy #2: TXA and Combined Hormonal Contraception



Caution ⚠️

Combined hormonal contraception is listed as contraindication to TXA in the United States³⁸

There is a theoretical concern that TXA will further increase VTE risk in patients taking combined hormonal contraception (CHC)⁶⁵




Only one published case reports **CHC + TXA** as a cause of thrombosis of a coronary vessel⁶⁸
→ This could have occurred with CHC alone or perhaps even without CHC...

Evidence that *may* suggest that CHC + TXA is safe:


Post-partum VTE risk:
~300/10,000 woman yrs⁶⁷
(due to hormonal effects similar to CHC)

WOMAN Trial⁵
In patients with PPH, there was **no increased risk of VTE with TXA**



Individualized care and shared decision making is required when considering TXA in combination with CHC for patients with heavy menstrual bleeding⁶⁵

- Counselling on individual risks of VTE
- Weigh **benefits of therapy** against **potential risk of thrombosis**




Did you know? 💡


CHC is **NOT** a risk factor for recurrent VTE while patients are anticoagulated⁶⁹

Studies on the use of CHC + TXA are needed to determine combined efficacy & safety
A scoping review⁷⁰ and ISTH registry are ongoing to address this controversy

Controversy #3: Tranexamic Acid and Hematuria



The concern is that TXA may encourage **clot formation in the ureter or renal pelvis** which many lead to **acute kidney injury**³⁷



One case report⁷¹ in a patient with PCKD presenting with **life-threatening hematuria**

- Bilateral nephrectomy avoided
- Clot obstructing ureters requiring **J stents**

Case report⁷² and case series⁷³ in patients with PCKD presenting with hematuria showed **evidence of benefit**

- Cessation of hematuria
- No acute kidney injury
- No thromboembolism

Research In Progress



*Systematic review of hematuria and acute renal failure with cyclokapron (TXA)*⁷⁴ **in progress** to better address this controversy

Larger studies likely needed to evaluate benefit and characterize risk of clot formation and acute kidney injury

TXA ILLUSTRATED REVIEW

Reference citations¹⁻⁷⁴

ACKNOWLEDGMENTS

Select images in this review were taken from <https://smart.servier.com> and are licensed under a Creative Commons Attribution 3.0 Unported License. Blood drop image (pages 1, 4-5, 9-11) was purchased from shutterstock (stock vector ID: 641207020)

RELATIONSHIP DISCLOSURE

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

NR and MS conceived and designed the manuscript. NR, NLJC, and MS wrote the paper.

ORCID

Nicole Relke <https://orcid.org/0000-0003-3416-6765>

TWITTER

Nicole Relke  @NicoleRelke

Nicholas L. J. Chornenki  @NicholasJC95

Michelle Sholzberg  @sholzberg

REFERENCES

- Watts G, Utako Okamoto (Obituary). *Lancet*. 2016;387:2286.
- Okamoto S, Okamoto U. Amino-methyl-cyclohexane-carboxylic acid: AMCHA. A new potent inhibitor of fibrinolysis. *Keio J Med*. 1962;11:105-115.
- Olldashi F, Kerçi M, Zhurda T, et al. Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet*. 2010;376:23-32.
- WHO Expert Committee. The selection and use of essential medicines (including the 17th WHO model list of essential medicines). *WHO Tech Rep Ser [Internet]*. 2011;2011:1-249.
- Shakur H, Roberts I, Fawole B, et al. Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with post-partum haemorrhage (WOMAN): an international, randomised, double-blind, placebo-controlled trial. *Lancet*. 2017;389:2105-2116.
- Thibault GE. Women in academic medicine. *Acad Med*. 2016;91:1045-1046.
- Mannucci PM. Hemostatic drugs. *NEJM*. 1998;339:245-253.
- Chapin JC, Hajjar KA. Fibrinolysis and the control of blood coagulation. *Blood Rev*. 2015;29:17-24.
- Lin H, Xu L, Yu S, Hong W, Huang M, Xu P. Therapeutics targeting the fibrinolytic system. *Exp Mol Med*. 2020;52:367-379.
- Linkins LA, Takach LS. Review of D-dimer testing: Good, Bad, and Ugly. *Int J Lab Hematol*. 2017;39:98-103.
- Johnson ED, Schell JC, Rodgers GM. The D-dimer assay. *Am J Hematol*. 2019;94:833-839.
- Tengborn L, Blombäck M, Berntorp E. Tranexamic acid - an old drug still going strong and making a revival. *Thromb Res*. 2015;135:231-242.
- Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924-926.
- Novikova N, Hofmeyr GJ, Cluver C. Tranexamic acid for preventing postpartum haemorrhage. *Cochrane Database Syst Rev*. 2015;1-62.
- Myles PS, Smith JA, Forbes A, et al. Tranexamic acid in patients undergoing coronary-artery surgery. *N Engl J Med*. 2017;376:136-148.
- Collaborators C-3 trial. Effects of tranexamic acid on death, disability, vascular occlusive events and other morbidities in patients with acute traumatic brain injury (CRASH-3): a randomised, placebo-controlled trial. *Lancet*. 2019;394:1713-1723.
- Ac B, Lethaby A, Farquhar C, Hickey M. Antifibrinolytics for heavy menstrual bleeding (Review). *Cochrane Database Syst Rev*. 2018;1-76.
- La S, Warner GT, Cope E. Treatment of menorrhagia with tranexamic acid. a double-blind trial summary: a double-blind trial. *Br Med J*. 1970;4:214-216.
- Lukes A, Moore K, Muse K, Gersten J. Tranexamic acid treatment for heavy menstrual bleeding. *J Obstet Gynecol*. 2010;116:865-867.
- Sprigg N, Flaherty K, Appleton JP, et al. Tranexamic acid for hyperacute primary IntraCerebral Haemorrhage (TICH-2): an international randomised, placebo-controlled, phase 3 superiority trial. *The Lancet*. 2018;391:2107-2115. [https://doi.org/10.1016/S0140-6736\(18\)31033-X](https://doi.org/10.1016/S0140-6736(18)31033-X)
- Wand O, Guber E, Guber A, Epstein Shochet G, Israeli-Shani L, Shitrit D. Inhaled tranexamic acid for hemoptysis treatment. *Chest*. 2018;154:1379-1384. <https://doi.org/10.1016/j.chest.2018.09.026>
- Calvo GS, De Granda-Orive I, Padilla DL. Inhaled tranexamic acid as an alternative for hemoptysis treatment. *Chest*. 2016;149:604. <https://doi.org/10.1016/j.chest.2015.10.016>
- Kagoma YK, Crowther MA, Douketis J, Bhandari M, Eikelboom J, Lim W. Use of antifibrinolytic therapy to reduce transfusion in patients undergoing orthopedic surgery: a systematic review of randomized trials. *Thromb Res*. 2009;123:687-696.
- Wang Z, Zhang H. Comparative effectiveness and safety of tranexamic acid plus diluted epinephrine to control blood loss during total hip arthroplasty: a meta-analysis. *J Orthop Surg Res*. 2018;13:1-11.
- Morimoto Y, Yoshioka A, Sugimoto M, Imai Y, Kirita T. Haemostatic management of intraoral bleeding in patients with von Willebrand disease. *Oral Dis*. 2005;11:243-248.
- Eghbali A, Melikof L, TaherAhmadi H, Bagheri B. Efficacy of tranexamic acid for the prevention of bleeding in patients with von Willebrand disease and Glanzmann thrombasthenia: a controlled, before and after trial. *Haemophilia*. 2016;22:e423-e426.
- Montroy J, Hutton B, Moodley P, et al. The efficacy and safety of topical tranexamic acid: a systematic review and meta-analysis. *Transfus Med Rev*. 2018;32:165-178.
- Zahed R, Mousavi Jazayeri MH, Naderi A, Naderpour Z, Saeedi M. Topical tranexamic acid compared with anterior nasal packing for treatment of epistaxis in patients taking antiplatelet drugs: randomized controlled trial. *Acad Emerg Med*. 2018;25:261-266.
- Joseph J, Martinez-Devesa P, Bellorini J, Burton MJ. Tranexamic acid for patients with nasal haemorrhage (epistaxis). *Cochrane Database Syst Rev*. 2018;2018:1-38.
- Geisthoff UW, Seyfert UT, Kübler M, Bieg B, Plinkert PK, König J. Treatment of epistaxis in hereditary hemorrhagic telangiectasia with tranexamic acid - a double-blind placebo-controlled cross-over phase IIIb study. *Thromb Res*. 2014;134:565-571.
- Shpilberg O, Blumenthal R, Sofer O, et al. A controlled trial of tranexamic acid therapy for the reduction of bleeding during treatment of acute myeloid leukemia. *Leuk Lymphoma*. 1995;19:141-144.
- Avvisati G, Büller HR, Ten CJW, Mandelli F. Tranexamic acid for control of haemorrhage in acute promyelocytic leukaemia. *Lancet*. 1989;334:122-124.
- Kim HJ, Moon SH, Cho SH, Lee JD, Kim HS. Efficacy and safety of tranexamic acid in melasma: a meta-analysis and systematic review. *Acta Derm Venereol*. 2017;97:776-781.
- Compels MM, Lock RJ. C1 inhibitor deficiency: management. *Clin Exp Dermatol*. 2005;30:737-740.
- Sentilhes L, Winer N, Azria E, et al. Tranexamic acid for the prevention of blood loss after vaginal delivery. *N Engl J Med*. 2018;379:731-742.
- Sentilhes L, Daniel V, Deneux-Tharoux C. TRAAP2-TRANexamic Acid for Preventing postpartum hemorrhage after cesarean delivery: a multicenter randomized, doubleblind, placebo-controlled trial-a study protocol. *BMC Pregnancy Childbirth*. 2020;20:1-11.
- Ker K, Roberts I, Chaudhri R, et al. Tranexamic acid for the prevention of postpartum bleeding in women with anaemia: study protocol for an international, randomised, double-blind, placebo-controlled trial. *Trials*. 2018;19:1-19.
- Pfizer Canada. *Product Monograph: Cyklokapron [Internet]*; 2018. <https://www.pfizer.ca/cyklokapron-tranexamic-acid> (accessed December 16, 2020).
- Tranexamic acid*. FDA. [Internet]; 2019. https://www.accessdata.fda.gov/drugsatfda_docs/label/2019/212020lbl.pdf (accessed January 25, 2021).
- Muse K, Lukes AS, Gersten J, Waldbaum A, Mabey RG, Trott E. Long-term evaluation of safety and health-related quality of life in women with heavy menstrual bleeding treated with oral tranexamic acid. *Women's Heal*. 2011;7:699-707.
- Lecker I, Wang DS, Whissell PD, Avramescu S, Mazer CD, Orser BA. Tranexamic acid-associated seizures: causes and treatment. *Ann Neurol*. 2016;79:18-26.

42. Gofton TE, Chu MWA, Norton L, et al. A prospective observational study of seizures after cardiac surgery using continuous EEG monitoring. *Neurocrit Care*. 2014;21:220-227.
43. Manji RA, Grocott HP, Leake J, et al. Seizures following cardiac surgery: the impact of tranexamic acid and other risk factors. *Can J Anesth*. 2012;59:6-13.
44. Kalavrouziotis D, Voisine P, Mohammadi S, Dionne S, Dagenais F. High-dose tranexamic acid is an independent predictor of early seizure after cardiopulmonary bypass. *Ann Thorac Surg [Internet]*. 2012;93:148-154. <https://doi.org/10.1016/j.athoracsur.2011.07.085>
45. Lin Z, Xiaoyi Z. Tranexamic acid-associated seizures: a meta-analysis. *Seizure [Internet]*. 2016;36:70-73. <https://doi.org/10.1016/j.seizure.2016.02.011>
46. Baharoglu MI, Germans MR, Rinkel GJE, et al. Antifibrinolytic therapy for aneurysmal subarachnoid haemorrhage. *Cochrane Database Syst Rev*. 2013;2013.
47. Imbesi S, Nettis E, Minciullo PL, et al. Hypersensitivity to tranexamic acid: a wide spectrum of adverse reactions. *Pharm World Sci*. 2010;32:416-419.
48. Li PH, Trigg C, Rutkowski R, Rutkowski K. Anaphylaxis to tranexamic acid—a rare reaction to a common drug. *J Allergy Clin Immunol Pract [Internet]*. 2017;5:839-841. <https://doi.org/10.1016/j.jaip.2016.12.014>
49. Connolly ES, Rabinstein AA, Carhuapoma JR, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2012;43:1711-1737.
50. Post R, Germans MR, Tjerkstra MA, et al. Ultra-early tranexamic acid after subarachnoid haemorrhage (ULTRA): a randomised controlled trial. *Lancet*. 2020;6736:1-7.
51. Hallberg L, Högdahl AM, Nilsson L, Rybo G. Menstrual blood loss—a population study. Variation at different ages and attempts to define normality. *Obstet Gynecol Surv*. 1967;22:653-654.
52. *Heavy menstrual bleeding clinical guideline 2007*. National Collaborating Centre for Women's and Children's Health. NICE; 2007.
53. Connell NT, Flood VH, Brignardello-Petersen R, et al. ASH ISTH NHF WFH 2021 guidelines on the management of von Willebrand disease. *Blood Adv*. 2021;5:301-325.
54. Kouides PA, Byams VR, Philipp CS, et al. Multisite management study of menorrhagia with abnormal laboratory haemostasis: a prospective crossover study of intranasal desmopressin and oral tranexamic acid. *Br J Haematol*. 2009;145:212-220.
55. Roberts I, Shakur-Still H, Afolabi A, et al. Effects of a high-dose 24-h infusion of tranexamic acid on death and thromboembolic events in patients with acute gastrointestinal bleeding (HALT-IT): an international randomised, double-blind, placebo-controlled trial. *Lancet*. 2020;395:1927-1936.
56. Roberts LN. Tranexamic acid in acute gastrointestinal bleeding – a cautionary tale. *J Thromb Haemost*. 2020;18:2440-2443.
57. Blasi A, Patel VC, Adelmeyer J, et al. Mixed fibrinolytic phenotypes in decompensated cirrhosis and acute-on-chronic liver failure with hypofibrinolysis in those with complications and poor survival. *Hepatology*. 2020;71:1381-1390.
58. Lisman T, De Groot PG, Meijers JCM, Rosendaal FR. Reduced plasma fibrinolytic potential is a risk factor for venous thrombosis. *Blood*. 2005;105:1102-1105.
59. Gernsheimer T, Brown SP, Triulzi DJ, et al. Effects of tranexamic acid prophylaxis on bleeding outcomes in hematologic malignancy: the a-TREAT trial. *Blood*. 2020;136:1-2.
60. Miller AB, Hoogstraten B, Staquet M, Winkler A. Reporting results of cancer treatment. *Cancer*. 1981;47:207-214.
61. Chornenki NLJ, Um KJ, Mendoza PA, et al. Risk of venous and arterial thrombosis in non-surgical patients receiving systemic tranexamic acid: a systematic review and meta-analysis. *Thromb Res [Internet]*. 2019;179:81-86. <https://doi.org/10.1016/j.thromres.2019.05.003>
62. Taeuber I, Weibel S, Herrmann E, et al. Association of intravenous tranexamic acid with thromboembolic events and mortality. A systematic review, meta-analysis, and meta-regression. *JAMA Surg*. 2021;156:1-14.
63. Ker K, Edwards P, Perel P, Shakur H, Roberts I. Effect of tranexamic acid on surgical bleeding: systematic review and cumulative meta-analysis. *BMJ*. 2012;344:1-13.
64. Ker K, Prieto-Merino D, Roberts I. Systematic review, meta-analysis and meta-regression of the effect of tranexamic acid on surgical blood loss. *Br J Surg*. 2013;100:1271-1279.
65. Thorne JG, James PD, Reid RL. Heavy menstrual bleeding: is tranexamic acid a safe adjunct to combined hormonal contraception? *Contraception [Internet]*. 2018;98:1-3. <https://doi.org/10.1016/j.contraception.2018.02.008>
66. Heinemann LAJ, Dinger JC. Range of published estimates of venous thromboembolism incidence in young women. *Contraception*. 2007;75:328-336.
67. Reid RL, Westhoff C, Mansour D, et al. Oral contraceptives and venous thromboembolism: consensus opinion from an international workshop held in Berlin, Germany in December 2009. *J Fam Plan Reprod Heal Care*. 2010;36:117-122.
68. Iacobellis G, Iacobellis G. Combined treatment with tranexamic acid and oral contraceptive pill causes coronary ulcerated plaque and acute myocardial infarction. *Cardiovasc Drugs Ther*. 2004;18:239-240.
69. Klok FA, Schreiber K, Stach K, et al. Oral contraception and menstrual bleeding during treatment of venous thromboembolism: expert opinion versus current practice. *Thromb Res*. 2017;153:101-107.
70. Meschino D, Sholzberg M. Evaluating the combined effect of anti-fibrinolytics and estrogen on the risk of thromboembolism: a scoping review. Available from: osf.io/edj28
71. Vujkovic B, Sabovic M. A successful treatment of life-threatening bleeding from polycystic kidneys with antifibrinolytic agent tranexamic acid. *Blood Coagul Fibrinolysis*. 2006;17:589-591.
72. AlAmeel T, West M. Tranexamic acid treatment of life-threatening hematuria in polycystic kidney disease. *Int J Nephrol*. 2011;2011:1-3.
73. Peces R, Aguilar A, Vega C. Medical therapy with tranexamic acid in autosomal dominant polycystic kidney disease patients with severe haematuria. *Nefrologia*. 2012;32:160-165.
74. Lee S, Fralick M, Sholzberg M, et al. Systematic-review of Hematuria and Acute Renal-failure with cycloKapron (SHARK); 2020.

How to cite this article: Relke N, Chornenki NLJ, Sholzberg M. Tranexamic acid evidence and controversies: An illustrated review. *Res Pract Thromb Haemost*. 2021;5:e12546. <https://doi.org/10.1002/rth2.12546>